
The Oscillation Length Resonance in the Transitions of Solar and Atmospheric Neutrinos Crossing the Earth Core

Serguey PETCOV **Scuola Internazionale Superiore di Studi Avanzati,
Istituto Nazionale di Fizica Nucleare, Sezione di Trieste,
I-34013 Trieste, Italy, and Bulgarian Academy of Sciences,
1784 Sofia, Bulgaria,
petcov@susy.sissa.it*

Abstract

The characteristic features of the neutrino oscillation length resonance, taking place in the $\nu_2 \rightarrow \nu_e$ and $\nu_\mu \rightarrow \nu_e$ ($\nu_e \rightarrow \nu_{\mu(\tau)}$) transitions in the Earth of the solar and atmospheric neutrinos crossing the Earth core, are reviewed. The resonance enhances dramatically these transitions at small mixing angles but differs from the MSW one. It can be responsible, e.g., for the small excess of e -like events at Zenith angles $\theta_z \sim (130^\circ - 180^\circ)$ and can produce at least part of the Zenith angle dependence of the rate of μ -like events, exhibited by the Super-Kamiokande atmospheric neutrino data.

1. Introduction

The $\nu_2 \rightarrow \nu_e$ and $\nu_\mu \rightarrow \nu_e$ ($\nu_e \rightarrow \nu_{\mu(\tau)}$) transitions of the solar and atmospheric neutrinos in the Earth, caused by neutrino mixing in vacuum ¹, are enhanced by a new type of resonance which differs from the MSW one and takes place when the neutrinos cross the Earth core [1]. The enhancement is particularly strong in the case of small mixing [1]. The conditions for existence of the new resonance include specific constraints on the neutrino oscillation length (and mixing angles) in the Earth mantle and core. When satisfied, these constraints

* Talk given at the Satellite Symposium “New Era in Neutrino Physics”, June 11 - 12, 1998, Tokyo Metropolitan University, Tokyo, Japan (to be published in the Proceedings of the Symposium).

1 The $\nu_2 \rightarrow \nu_e$ transition probability accounts, as is well-known, for the Earth effect in the solar neutrino MSW survival probability in the case of the two-neutrino $\nu_e \rightarrow \nu_{\mu(\tau)}$ and $\nu_e \rightarrow \nu_s$ transition solutions of the solar neutrino problem, ν_s being a sterile neutrino.

ensure that certain oscillating factors in the probabilities of the $\nu_2 \rightarrow \nu_e$ and $\nu_\mu \rightarrow \nu_e$ ($\nu_e \rightarrow \nu_{\mu(\tau)}$) transitions, P_{e2} and $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$, have maximal values and that this produces a resonance maximum in P_{e2} and $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$. Correspondingly, the term “neutrino oscillation length resonance” (NOLR) or simply “oscillation length resonance” was used in [1] to denote the resonance of interest. The NOLR is in many respects similar, e.g., to the resonance transforming a circularly polarized LH photon into a RH one when the photon traverses three layers of optically active medium such that the optical activity and the widths of the first and third layers are identical but differ from those of the second layer ².

The present article contains a brief review of the specific features of the oscillation length resonance in the simplest case of two-neutrino transitions of the Earth core crossing solar and atmospheric neutrinos. We discuss also the implications of the NOLR enhancement of these transitions [1,2,3,4] for the interpretation of the results of the Super-Kamiokande experiment [5,6].

2. The Neutrino Oscillation Length Resonance (NOLR) in the Transitions of Solar and Atmospheric Neutrinos Traversing the Earth

According to the existing Earth models [7], the matter density and the chemical composition change mildly within the two main density structures in the Earth - the mantle and the core. As a consequence, all the interesting features of the solar and atmospheric neutrino transitions in the Earth, including those related to the neutrino oscillation length resonance, can be understood quantitatively assuming that the neutrinos crossing the mantle and the core traverse effectively two layers with constant but different densities, $\bar{\rho}_{man}$ and $\bar{\rho}_c$, and chemical composition or electron fraction numbers, Y_e^{man} and Y_e^c [1,3,8,9]. The neutrino transitions of interest in the two-layer model of the Earth density distribution ³ result from two-neutrino oscillations taking place i) first in the mantle over a distance X' with a mixing angle θ'_m and oscillation length L_{man} , ii) then in the core over a distance X'' with different mixing angle θ''_m and oscillation length L_c , and iii) again in the mantle over a distance X' with θ'_m and L_{man} . Because $\bar{\rho}_{man,c} \neq 0$, $\bar{\rho}_{man} \neq \bar{\rho}_c$ and $Y_e^{man} \sim Y_e^c$, and due to the matter effect, we have for given neutrino energy E , mass squared difference $\Delta m^2 > 0$ and vacuum mixing angle θ , $\cos 2\theta > 0$ (see, e.g., [11]): $\theta'_m, \theta''_m \neq \theta$, $\theta'_m \neq \theta''_m$, $L_{man} \neq L_c$ and $L_{man,c} \neq L^{vac}$, $L^{vac} = 4\pi E / \Delta m^2$ being the neutrino oscillation length in vacuum.

² There exists also a beautiful analogy between the NOLR and the electron spin-flip resonance realized in a specific configuration of magnetic fields (see [1] for a more detailed discussion).

³ The densities $\bar{\rho}_{man,c}$ should be considered as mean densities along the neutrino trajectories. In the Earth models [7] one has: $\bar{\rho}_{man} \cong (4 - 5) \text{ g/cm}^3$ and $\bar{\rho}_c \cong (11 - 12) \text{ g/cm}^3$. For Y_e one can use the standard values [7,10] (see also [4]) $Y_e^{man} = 0.49$ and $Y_e^c = 0.467$.

For fixed X' and X'' the oscillation length resonance occurs [1] if i) the relative phases acquired by the energy eigenstate neutrinos in the mantle and in the core, $\Delta E' X' = 2\pi X'/L_{man}$ and $\Delta E'' X'' = 2\pi X''/L_c$, are correlated,

$$\Delta E' X' = \pi(2k+1), \quad \Delta E'' X'' = \pi(2k'+1), \quad k, k' = 0, 1, 2, \dots, \quad (1)$$

and if ii) the inequalities

$$\sin^2(2\theta''_m - 4\theta'_m + \theta) - \sin^2 \theta > 0, \quad (2a)$$

$$\begin{aligned} & \sin^2(2\theta''_m - 4\theta'_m + \theta) \sin(2\theta''_m - 2\theta'_m) \sin(2\theta'_m - \theta) \cos(2\theta''_m - 4\theta'_m + \theta) \\ & + \frac{1}{4} \sin^2 \theta \sin(4\theta''_m - 8\theta'_m + 2\theta) \sin(4\theta''_m - 4\theta'_m) < 0. \end{aligned} \quad (2b)$$

are fulfilled. Conditions (1) are universal, i.e., they are valid for P_{e2} , $P(\nu_\mu \rightarrow \nu_e) = P(\nu_e \rightarrow \nu_{\mu(\tau)})$, etc. while (2a) and (2b) correspond only to P_{e2} [1]. When equalities (1) hold, (2a) and (2b) ensure that P_{e2} has a maximum. In the region of the NOLR maximum, e.g., for $\Delta E' X' \cong \pi(2k+1)$, P_{e2} is given by [1]:

$$P_{e2} \cong \sin^2 \theta + \frac{1}{4} [1 - \cos \Delta E'' X''] [\sin^2(2\theta''_m - 4\theta'_m + \theta) - \sin^2 \theta]. \quad (3)$$

At the NOLR maximum we have [1] (Fig. 1):

$$P_{e2}^{max} = \sin^2(2\theta''_m - 4\theta'_m + \theta). \quad (4)$$

The analogs of eqs. (2) - (4) for $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$ can be obtained by formally setting $\theta = 0$ while keeping $\theta'_m, \theta''_m \neq 0$ in eqs. (2) - (4): thus, (2a) becomes $\sin^2(2\theta''_m - 4\theta'_m) > 0$ and is always fulfilled, while (2b) transforms into [1]

$$\cos(2\theta''_m - 4\theta'_m) < 0. \quad (5)$$

Note that since X' and X'' are fixed for a given neutrino trajectory, which is specified by its Nadir angle h , the NOLR conditions (1) are constraints on the oscillation lengths L_{man} and L_c . At $\sin^2 2\theta \lesssim 0.02$ one of the two requirements in (1) is equivalent for $k = k' = 0$ to the physical condition [1]

$$\pi \left(\frac{1}{X'} + \frac{1}{X''} \right) \cong \sqrt{2} G_F \frac{1}{m_N} (Y_e^c \bar{\rho}_c - Y_e^{man} \bar{\rho}_{man}).$$

Remarkably enough, for the $\nu_2 \rightarrow \nu_e$ and $\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)}$ transitions of the solar and atmospheric neutrinos in the Earth, the NOLR conditions (1) with $k = k' = 0$ are approximately fulfilled at small $\sin^2 2\theta$ in the regions where (2a) - (2b) or (5) hold [1]. The associated NOLR maxima in P_{e2} and $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$ are absolute maxima and dominate in P_{e2} and $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$ [1,3] (Fig. 2): the




Fig. 1. The dependence of the function $\sin^2(2\theta''_m - 4\theta'_m + \theta)$, eq. (4), on $\rho_r = 1.32 \times 10^7 \Delta m^2 [eV^2] \cos 2\theta / E [MeV] \text{ g/cm}^3 \approx \rho_{man}^{res}$, ρ_{man}^{res} being the resonance density in the Earth mantle [1], for $\sin^2 2\theta = 0.01$ (from [3]).

values of the probabilities at these maxima in the case of two-neutrino mixing are considerably larger - by a factor of $\sim (2.5 - 4.0)$ ($\sim (3.0 - 7.0)$), than the values of P_{e2} and $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$ at the local maxima associated with the MSW effect in the Earth core (mantle). The magnitude of the NOLR enhancement depends on the neutrino trajectory through the Earth core: the enhancement is maximal for the center-crossing neutrinos [1,3]. Even at small mixing angles the resonance is relatively wide both in the neutrino energy and the Nadir angle [1,3] (Fig. 2). The NOLR (relative) energy width is given for $\sin^2 2\theta < 0.05$ approximately by

$$\frac{\Delta E}{E_{max}} \cong \left(2 \left(1 + \frac{\sqrt{2}}{\pi} \frac{1}{m_N} G_F Y_e^{man} \bar{\rho}_{man} X' \right) \right)^{-1}. \quad (6)$$

It practically does not depend on $\sin^2 2\theta$ and varies slowly with the change of h : for $h \lesssim 25^\circ$ one has ⁴ $\Delta E / E_{max} \cong 0.30$.

Let us note that the dominating maximum due to the NOLR can *now* be recognized in certain plots of the probabilities P_{e2} and/or $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$, shown in some of the articles discussing the Earth effect in the transitions of solar and atmospheric neutrinos (see, e.g., [12,13]). This maximum was invariably interpreted to be due to the MSW effect in the Earth core ⁵ before the study [1].

⁴ For further details and a study of the NOLR effects in the transitions $\nu_2 \rightarrow \nu_e$ and $\nu_e \rightarrow \nu_s$ ($\nu_e - \nu_s$ mixing), and in $\bar{\nu}_\mu \rightarrow \bar{\nu}_s$ (or $\nu_\mu \rightarrow \nu_s$) at small mixing angles, see [1,3].

⁵ Let us add also that the analysis performed in [1] and summarized above differs substantially from the studies [14] of the resonance enhancement of the $\nu_e \rightarrow \nu_{\mu(\tau)}$ transitions in matter varying *periodically* along the neutrino path (parametric resonance). The density change along the path of a neutrino crossing the Earth core is not periodic (even in the two-layer




Fig. 2. The probability $P(\nu_{e(\mu)} \rightarrow \nu_{\mu;\tau(e)})$ as a function of h and $E/\Delta m^2$ for $\sin^2 2\theta = 0.01$ [3]. The NOLR absolute maximum for $h \approx (0^\circ - 28^\circ)$ is clearly seen at $E/\Delta m^2 \approx (1.3 - 1.6) \times 10^6$ MeV/eV². The local maximum at $E/\Delta m^2 \sim 2.5 \times 10^6$ MeV/eV² is due to the MSW effect in the Earth mantle.

Finally, in [15] the $\nu_\mu \rightarrow \nu_s$ transitions in the Earth were considered for $\sin^2 2\theta \cong 1$. It was noticed that in the region where $\sqrt{2}G_F \bar{N}_n^{man,c} \gg \Delta m^2/E$, $\bar{N}_n^{man,c}$ being the neutron number density, a new maximum in $P(\nu_\mu \rightarrow \nu_s)$ appears when $\sqrt{2}G_F \bar{N}_n^{man(c)} X^{(n)} \cong 2\pi$, found to hold at $h \sim 28^\circ$. The height of the maximum is comparable to the heights of the other “ordinary” maxima present in $P(\nu_\mu \rightarrow \nu_s)$ for $\sin^2 2\theta \cong 1$. It is stated in [15] that the effect does not take place in the $\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)}$ transitions and the case of solar neutrinos is not discussed.

3. The Super-Kamiokande Atmospheric Neutrino Data and the Neutrino Oscillation Length Resonance

The implications of the NOLR enhancement of the probability P_{e2} for the tests of the MSW $\nu_e \rightarrow \nu_{\mu(\tau)}$ and $\nu_e \rightarrow \nu_s$ solutions of the solar neutrino problem are discussed in detail in refs. [1,2,3,4]. In the case of the small mixing angle (nonadiabatic) $\nu_e \rightarrow \nu_{\mu(\tau)}$ solution, the NOLR produces [4] a much bigger - by a factor of ~ 6 , day-night asymmetry in the Super-Kamiokande sample of solar neutrino events, whose night fraction is due to the *core-crossing* neutrinos, in

model it falls short of making one and a half periods) and the results obtained in [14] do not apply to the case of transitions of the Earth core crossing neutrinos, studied in [1].

comparison with the asymmetry determined by using the *whole night* event sample. The Super-Kamiokande collaboration has successfully exploited this result [3] in the analysis of their solar neutrino data in terms of the MSW effect [5].

The NOLR enhancement of the $\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)}$ transitions of the atmospheric neutrinos crossing the Earth has important implications for the interpretation of the Super-Kamiokande atmospheric neutrino data as well [1]. The Zenith angle dependence of the rates of sub-GeV and multi-GeV μ -like events, observed in the Super-Kamiokande experiment, provides a strong evidence for oscillations of the atmospheric ν_{μ} ($\bar{\nu}_{\mu}$) [6]. The data is best described in the case of two-neutrino mixing by $\nu_{\mu}(\bar{\nu}_{\mu}) \leftrightarrow \nu_{\tau}(\bar{\nu}_{\tau})$ vacuum oscillations with parameters $\Delta m^2 \cong (0.5 - 6.0) \times 10^{-3} \text{ eV}^2$ and $\sin^2 2\theta \cong (0.8 - 1.0)$ [6].

It is quite remarkable that for values of $\Delta m^2 \sim (0.5 - 6.0) \times 10^{-3} \text{ eV}^2$ suggested by the Super-Kamiokande data, the NOLR strongly enhances the $\nu_{\mu} \rightarrow \nu_e$ (and $\nu_e \rightarrow \nu_{\mu(\tau)}$) transitions of the atmospheric neutrinos crossing the Earth core, making the transitions detectable even at small mixing angles [1]. Indeed, for, e.g., $\sin^2 2\theta = 0.01$, $\Delta m^2 = 0.5; 1.0; 5.0 \times 10^{-3} \text{ eV}^2$, and $h = 0^0$ (Earth center crossing), the absolute maximum in $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$ due to the NOLR takes place at $E \cong 0.75; 1.50; 7.5 \text{ GeV}$. The ν_e and ν_{μ} with such energies contribute either to the sub-GeV or to the multi-GeV e -like and μ -like Super-Kamiokande event samples [6]. At small mixing angles the NOLR enhancement holds practically for all neutrino trajectories through the Earth core [1,3]. Accordingly, it was suggested in [1,2] (see also [3]) that, depending on the value of Δm^2 , the small excess of e -like events in the region $-1 \leq \cos \theta_z \leq -0.6$, $\theta_z = \pi - h$ being the Zenith angle, either in the sub-GeV or in the multi-GeV sample, observed (in both samples) in the Super-Kamiokande experiment [6], is due to $\nu_{\mu} \rightarrow \nu_e$ small mixing transitions, $\sin^2 2\theta_{e\mu} \cong (0.01 - 0.10)$, strongly enhanced by the NOLR as neutrinos cross the Earth core on the way to the detector. The same NOLR enhanced transitions should produce at least part of the zenith angle dependence, exhibited by the Super-Kamiokande sub-GeV and multi-GeV μ -like data [2,3].

The transitions of interest arise naturally, e.g., in a three-neutrino mixing scheme, in which the small mixing angle MSW $\nu_e \rightarrow \nu_{\mu}$ transitions with $\Delta m_{21}^2 \sim (4 - 8) \times 10^{-6} \text{ eV}^2$, or large mixing angle $\nu_e \leftrightarrow \nu_{\mu}$ oscillations with $\Delta m_{21}^2 \sim 10^{-10} \text{ eV}^2$, provide the solution of the solar neutrino problem and the atmospheric neutrino anomaly is due to $\nu_{\mu} \leftrightarrow \nu_{\tau}$ oscillations with $\Delta m_{31}^2 \sim (0.5 - 6.0) \times 10^{-3} \text{ eV}^2$ [1]. For $\Delta m_{31}^2 \gg \Delta m_{21}^2$ the three-neutrino $\nu_{\mu} \rightarrow \nu_e$ and $\nu_e \rightarrow \nu_{\mu(\tau)}$ transition probabilities reduce to a two-neutrino transition probability equal to $P(\nu_e \rightarrow \nu_{\tau})$ [16], with Δm_{31}^2 and $\sin^2 2\theta_{13} = 4|U_{e3}|^2(1 - |U_{e3}|^2)$ playing the role of the two-neutrino oscillation parameters⁶, $\theta_{13} \equiv \theta_{e\mu}$, where U_{e3} is the $e - \nu_3$ element of the

6 Analytic expression for the probability $P(\nu_e \rightarrow \nu_{\tau})$ can be found in [1] (see also eq. (3)).

lepton mixing matrix U . The data [6,17] imply: $\sin^2 2\theta_{13} \lesssim 0.20$ ⁷. The fluxes of atmospheric $\nu_{e,\mu}$ with energy E , crossing the Earth along a trajectory with Zenith angle θ_z before reaching the detector, $\Phi_{\nu_{e,\mu}}$, are given in this scheme by [2,3,16]:

$$\Phi_{\nu_e} \cong \Phi_{\nu_e}^0 \left(1 + [s_{23}^2 r(E, \theta_z) - 1] P(\nu_e \rightarrow \nu_\tau) \right), \quad (7)$$

$$\begin{aligned} \Phi_{\nu_\mu} \cong \Phi_{\nu_\mu}^0 (1 + s_{23}^4 [(s_{23}^2 r(E, \theta_z))^{-1} - 1] P(\nu_e \rightarrow \nu_\tau) \\ - 2c_{23}^2 s_{23}^2 [1 - \text{Re}(e^{-i\kappa} A(\nu_\tau \rightarrow \nu_\tau))]), \end{aligned} \quad (8)$$

where $\Phi_{\nu_{e(\mu)}}^0 = \Phi_{\nu_{e(\mu)}}^0(E, \theta_z)$ is the $\nu_{e(\mu)}$ flux in the absence of neutrino transitions/oscillations, $s_{23}^2 \equiv |U_{\mu 3}|^2 / (1 - |U_{e 3}|^2) \leq 1$, $U_{\mu 3}$ being the $\mu - \nu_3$ element of U , $c_{23}^2 = 1 - s_{23}^2$, and $r(E, \theta_z) \equiv \Phi_{\nu_\mu}^0 / \Phi_{\nu_e}^0$. The two-neutrino transition probability amplitude $A(\nu_\tau \rightarrow \nu_\tau)$ is given in the two-layer model of the Earth density distribution by eq. (1) in ref. [1] in which first θ is formally set to $\pi/2$ while θ'_m and θ''_m are kept arbitrary, and after that $\sin \theta'_m$ ($\sin \theta''_m$) and $\cos \theta'_m$ ($\cos \theta''_m$) are interchanged. The phase κ has the form [2,3,16]:

$$\begin{aligned} \kappa \cong \frac{1}{2} \left[\frac{\Delta m_{31}^2}{2E} X + \sqrt{2} G_F \frac{1}{m_N} (X'' Y_e^c \bar{\rho}_c + 2X' Y_e^{man} \bar{\rho}_{man}) - 2\Delta E' X' - \Delta E'' X'' \right] \\ - \frac{\Delta m_{21}^2}{2E} X \cos 2\varphi_{12}, \end{aligned} \quad (9)$$

where $X = X'' + 2X'$ and $\cos 2\varphi_{12} = (|U_{e1}|^2 - |U_{e2}|^2) / (1 - |U_{e3}|^2)$. The interpretation of the Super-Kamiokande data [6] in terms of $\nu_\mu \leftrightarrow \nu_\tau$ oscillations requires $s_{23}^2 \cong (0.30 - 0.70)$, with 0.5 being the statistically preferred value. For the predicted ratio $r(E, \theta_z)$ of the ν_μ and ν_e fluxes for $-1 \leq \cos \theta_z \lesssim -0.8$ one has [19,20]: $r(E, \theta_z) \cong (2.0 - 2.5)$ for the neutrinos contributing to the sub-GeV event samples, and $r(E, \theta_z) \cong (2.6 - 4.5)$ for those giving the main contribution to the multi-GeV samples. If $s_{23}^2 = 0.5$ and $r(E, \theta_z) \cong 2.0$ we have $(s_{23}^2 r(E, \theta_z) - 1) \cong 0$ and the possible effects of the $\nu_\mu \rightarrow \nu_e$ and $\nu_e \rightarrow \nu_{\mu(\tau)}$ transitions on the ν_e flux and correspondingly on the sub-GeV e -like event sample, would be strongly suppressed even if these transitions are maximally enhanced by the NOLR. The indicated suppression may actually be taking place⁸. Note, however, that the factor $(s_{23}^2 r(E, \theta_z) - 1)$ can be as large as ~ 0.6 for the sub-GeV neutrinos and that the effects of the NOLR enhanced $\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)}$ transitions in the sub-GeV e -like event sample can be suppressed by the specific way this sample is selected

⁷ Thus, searching for the $\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)}$ transitions of the atmospheric neutrinos, amplified by the NOLR, can provide also unique information about the magnitude of U_{e3} [18].

⁸ Indeed, a more detailed investigation [18] performed within the indicated three-neutrino mixing scheme reveals, in particular, that the excess of e -like events in the Super-Kamiokande sub-GeV data at $-1 \leq \cos \theta_z \leq -0.6$ seems unlikely to be due to small mixing angle, $\sin^2 2\theta_{13} \lesssim 0.20$, $\nu_\mu \rightarrow \nu_e$ transitions amplified by the oscillation length resonance.

from the data in the Super-Kamiokande experiment [18]. For the multi-GeV neutrinos we have $(s_{23}^2 r(E, \theta_z) - 1) \gtrsim 0.3$ (0.9) for $s_{23}^2 = 0.5$ (0.7). The factor $(s_{23}^2 r(E, \theta_z) - 1)$ amplifies the effect of the $\nu_\mu \rightarrow \nu_e$ transitions in the e -like sample for $E \gtrsim (5 - 6)$ GeV, for which $r(E, \theta_z) \gtrsim 4$ [19,20]. This discussion suggests that the effects of the neutrino oscillation length resonance in the Super-Kamiokande e -like multi-GeV data for $\Delta m_{31}^2 \sim (2 - 6) \times 10^{-3} \text{ eV}^2$ should be much larger than those in the e -like sub-GeV data for $\Delta m_{31}^2 \sim (0.5 - 1.0) \times 10^{-3} \text{ eV}^2$ [2,3,18]. As similar analysis of the expression (8) for Φ_{ν_μ} shows, they are also expected to be larger (within the 3- ν mixing scheme considered) than the NOLR effects in the multi-GeV μ -like event sample [2,3,18]. In all samples, except, perhaps, the sub-GeV e -like one, the NOLR effects can be observable.

4. Conclusions

The neutrino oscillation length resonance should be present in the $\nu_2 \rightarrow \nu_e$ transitions of the solar neutrinos crossing the Earth core on the way to the detector, if the solar neutrino problem is due to small mixing angle MSW $\nu_e \rightarrow \nu_\mu$ transitions in the Sun. The same resonance should be operative also in the $\nu_\mu \rightarrow \nu_e$ ($\nu_e \rightarrow \nu_{\mu(\tau)}$) small mixing angle transitions of the atmospheric neutrinos crossing the Earth core if the atmospheric ν_μ and $\bar{\nu}_\mu$ indeed take part in large mixing vacuum $\nu_\mu(\bar{\nu}_\mu) \leftrightarrow \nu_\tau(\bar{\nu}_\tau)$ oscillations with $\Delta m^2 \sim (0.5 - 6.0) \times 10^{-3} \text{ eV}^2$, as is strongly suggested by the Super-Kamiokande data [6], and if all three flavour neutrinos are mixed in vacuum. Actually, the resonance may have already manifested itself in the excess of e -like events at $-1 \leq \cos \theta_z \leq -0.6$ observed in the Super-Kamiokande multi-GeV atmospheric neutrino data [1,2,3,18]. And it can be responsible for at least part of the strong zenith angle dependence present in the Super-Kamiokande multi-GeV and sub-GeV μ -like data [2,3,18].

5. Acknowledgements.

The author would like to thank H. Minakata and O. Yasuda for their kind hospitality during the *New Era in Neutrino Physics* Symposium.

6. References

1. S.T. Petcov, Phys. Lett. **B434** (1998) 321.
2. S.T. Petcov, Talk given at the Workshop “Neutrino Physics and Astrophysics: from Solar to Ultra-High Energy”, Aspen Center for Physics, June 29 - July 12, 1998, Aspen, U.S.A.
3. M. Chizhov, M. Maris and S.T. Petcov, Report SISSA 53/98/EP, 31 July 1998 (hep-ph/9810501); see also: S.T. Petcov, hep-ph/9809587.
4. M. Maris and S.T. Petcov, Phys. Rev. **D56** (1997) 7444, and hep-ph/9803244.

5. M. Nakahata (Super-Kamiokande Collaboration), these Proceedings.
6. T. Kajita, (Super-Kamiokande Collaboration), these Proceedings.
7. F.D. Stacey, *Physics of the Earth*, John Wiley and Sons, New York, 1977; A.D. Dziewonski and D.L. Anderson, Phys. Earth Planet. Interiors **25** (1981) 297.
8. P.I. Krastev and S.T. Petcov, Phys. Lett. **B205** (1988) 84.
9. M. Maris and S.T. Petcov, in preparation; M. Maris, Q.Y. Liu and S.T. Petcov, study performed in December of 1996 (unpublished).
10. R. Jeanloz, Annu. Rev. Earth Planet. Sci. **18** (1990) 356; C.J. Allègre et al., Earth Planet. Sci. Lett. **134** (1995) 515.
11. S.T. Petcov, Lecture Notes in Physics, vol. **512** (eds. H. Gausterer and C.B. Lang, Springer, 1998), p. 281.
12. S.P. Mikheyev and A.Yu. Smirnov, Proc. of the Moriond Workshop on Massive Neutrinos, 1986 (eds. O. Fackler and J. Tran Thanh Van, Editions Frontières, France, 1986), p. 355; E.D. Carlson, Phys. Rev. **D34** (1986) 1454; A. Dar et al., Phys. Rev. **D35** (1987) 3607; M. Cribier et al., Phys. Lett. **B182** (1986) 89; A.J. Baltz and J. Weneser, Phys. Rev. **D35** (1987) 528; J.M. Gelb, W. Kwong and S.P. Rosen, Phys. Rev. Lett. **78** (1997) 2296.
13. Q.Y. Liu, M. Maris and S.T. Petcov, Phys. Rev. **D56** (1997) 5991.
14. V.K. Ermilova et al., Short Notices of the Lebedev Institute **5** (1986) 26; E.Kh. Akhmedov, Yad. Fiz. **47** (1988) 475; P.I. Krastev and A.Yu. Smirnov, Phys. Lett. **B226** (1989) 341.
15. Q.Y. Liu and A.Yu. Smirnov, hep-ph/9712493.
16. S.T. Petcov, Phys. Lett. **B214** (1988) 259.
17. M. Appolonio et al. (CHOOZ Collaboration), Phys. Lett. **B420** (1998) 397.
18. S.T. Petcov, L. Wolfenstein and O. Yasuda, work in progress.
19. M. Honda et al., Phys. Rev. **D52** (1995) 4985.
20. V. Agraval et al., **D53** (1996) 1314.